arXiv:1207.5477v1 [nucl-th] 23 Jul 2012

Shell model description of odd sulfur isotopes

P.C. Srivastava,[1,*] Jorge G. Hirsch,[1] M.J. Ermamatov,[1,2] and V.K.B. Kota[3]

[1]*Instituto de Ciencias Nucleares, Universidad Nacional*
Autónoma de México, 04510 México, D.F., Mexico
[2]*Institute of Nuclear Physics, Ulughbek, Tashkent 100214, Uzbekistan*
[3]*Physical Research Laboratory, Ahmedabad 380 009, India*

(Dated: July 24, 2012)

Abstract

The low energy structure of ^{35,37,39,41}S isotopes is described performing shell model calculations with the recently devised SDPF-U interaction. Protons and neutrons are restricted to the *sd*-shell for $N < 20$, neutrons start filling the fp-shell for $N > 20$. Natural parity states are described with only in-shell mixing, opposite parity states with 1p-1h inter-shell neutron excitations. Energy levels and intensities of electromagnetic transitions are compared with experimental data when available.

PACS numbers: 21.60.Cs; 27.30.+t ; 27.40.+z

---

[*] E-mail: praveen.srivastava@nucleares.unam.mx

I. INTRODUCTION

The coexistence of normal and intruder, sometimes deformed, configurations in the low energy region [1] around the shell gaps $N = 20$ and $N = 28$, and the evidence of the erosion of these gaps [2], has been the focus of recent experimental investigations on Mg ($Z = 12$) [1], Si ($Z = 14$) [3], S ($Z = 16$) [4, 5] and Ar ($Z = 18$) isotopes [6]. The reaction analysis by Fortune [7] on $^{32}_{12}\mathrm{Mg}_{20}$ reveals that the 0^+ ground state has a normal (0p-0h) configuration, while the excited 0^+ state is an intruder (2p-2h). A strongly deformed intruder ground state has been reported in ^{31}Mg by the Leuven group [8, 9]. For ^{33}Mg, the parity of the spin $I = 3/2$ ground-state remains a puzzle [10–13]. For this region, a reduction in shell gap for the $s_{1/2}$ and $d_{3/2}$ proton orbitals has been reported for P, Cl and K isotopes [14–16]. The particle-hole excitations involving intruder states play an important role in the study of nuclei lying between *island of inversion* and the valley of stability. In Ar, the development of collectivity near the $Z = 28$ shell gap has been reported in [17], where nonaxially symmetric deformation was assigned to ^{48}Ar. In analogy to ^{32}Mg, with $N = 20$, a new *island of inversion* around $N = 40$ has been predicted in Ref. [18].

Sulfur isotopes exhibit many interesting properties: ^{40}S and ^{42}S are deformed, ^{41}S exhibit γ-soft properties [19] and collectivity [20], shape coexistence is found in ^{43}S [21], shape and configuration coexistence in ^{44}S [4, 22, 23]. Triple configuration coexistence is a topic of current research [24]. Experimental data are indicating an erosion of $N = 28$ shell gap in the ^{44}S and ^{42}Si isotones [3, 25].

Fig. 1 shows the systematics for $B(E2)$ values (top) and the energies of the $E(2^+_1)$ and $E(4^+_1)$ states (bottom), as a function of the neutron number, for even-even sulfur isotopes. A clear depletion in the $B(E2)$ value and a peak in the excitation energies is observed at ^{36}S with N = 20. As the neutron number increases, the $B(E2)$ values also grow up, and the excitation energies assume nearly constant low values. These two observables are associated with the development of collectivity. Sulfur isotopes have also an astrophysical importance. In the nuclear synthesis of heavy Ca-Ti-Cr isotopes, neutron-rich sulfur isotopes play an important role [26].

From the theoretical side, Kaneko *et al.* [27], recently reported calculations for positive parity states along the even-even sulfur chain using extended pairing plus quadrupole-quadrupole forces including monopole interactions (EPQQM). Monte Carlo shell model calculation based



FIG. 1. (Color online) Experimental $B(E2)$ values (top) and $E(2_1^+)$ and $E(4_1^+)$ energies (bottom) for even sulfur isotopes.

on the QMCD method describing oxygen to silicon with N $\sim$ 20 has been reported in [28]. We have performed shell-model (SM) studies in the fp shell for Fe, Mn, and Co isotopes [29–32], in the fpg shell nuclei for $^{71-79}$Ga [33], Ni, Cu and Zn isotopes [34, 35].

In this paper, the reported low energy states in ^{35,37,39,41}S isotopes, with both positive and negative parity, are interpreted in the frame work of shell model in $sd-pf$ space. While intruder states in the full $psdpf$ model space with a ^{4}He core allowing one nucleon jump between the major shells have been described in Ref. [36] with a new PSDPF interaction, to the best of our knowledge a detailed analysis of opposite parity band in these four odd-mass nuclei is presented for the first time here.

The paper is arranged as follows: in Sec. II, we discuss the effective interactions which are used in this work and provide some details of calculations. Results of calculations for the energy levels and electromagnetic transition probabilities of the ^{35,37,39,41}S isotopes are prresented in Sec. III. Finally, conclusions are drawn in Sec. IV.

II. EFFECTIVE INTERACTIONS

We present SM calculations in the $sd-pf$ space, using SDPF-U [37] interaction. This interaction is designed to be used in the valence space spanned by the orbitals $0d_{5/2}$, $1s_{1/2}$, $0d_{3/2}$, $0f_{7/2}$, $0f_{5/2}$, $1p_{3/2}$ and $1p_{1/2}$, for both protons and neutrons. The single-particle energies employed for the $0d_{5/2}$, $1s_{1/2}$, $0d_{3/2}$, $0f_{7/2}$, $1p_{3/2}$, $0f_{5/2}$ and $1p_{1/2}$ orbitals are -3.699,

-2.915, +1.895, +6.220, +6.314, +11.450, +6.479 MeV, respectively.

III. RESULTS AND DISCUSSION

A comparison of the calculated positive- and negative-parity states with the experimental data for the odd isotopes of sulfur is presented in Figs. 2-5. Experimental information for the energy levels in odd sulfur isotopes is available for ^{35}S, ^{37}S and ^{41}S isotopes [39]. Excited levels of ^{39}S have been measured at Legnaro National Laboratory (LNL) [40].

Protons and neutrons are restricted to the sd-shell for N $<$ 20, neutrons start filling the fp-shell for N $>$ 20. Natural parity states are described with only in-shell mixing, opposite parity states with 1p-1h inter-shell neutron excitations.

For the positive-parity levels in ^{35}S we perform 0p-0h calculations, i.e. both protons and neutrons are only allowed to occupy the sd shell. Negative-parity levels can be obtained for these nuclei exciting one neutron from the sd shell to the pf shell. The parity of the ground state becomes negative when reaching ^{37}S, since one or more neutrons already occupy the pf shell. Now negative-parity levels are obtained by 0p-0h and 1p-1h calculations give the positive parity bands.

The angular momentum and parity of the lowest energy states with both parities can be given a simple qualitative interpretation, associated with the last uncoupled neutron. In ^{35}S, $N = 19$, it occupies the $d_{3/2}$ orbital. For $N = 21$ and 23 the $f_{7/2}$ orbital is filled, and the ground state has $J = 7/2$ and negative parity. As in many other chains of isotopes, the energy of the excited state $J = 7/2^-$ is high in the lighter isotopes, and decreases for increasing N until it become the ground state, while the $J = 3/2^+$ becomes the first excited state with opposite parity, whose energy moves up. The subtle effects associated with configuration mixing and the role of the residual interaction can be observed following the gap between the $J = 7/2^-$ and $J = 3/2^-$ states. In ^{35}S they are the first and second excited states of negative parity, always in this order. In ^{37}S the ground state has $J = 7/2^-$ and the first excited state $J = 3/2^-$, but the theoretical predictions put both levels at near degeneracy in ^{39}S, and assign $J = 5/2^-$ to the ground state in ^{41}S, with the $J = 7/2^-$ the first excited state, in agreement with the preliminar experimental information. This type of behavior has been reported in [42], where a shell model study for phosphorous odd-odd isotopes was performed using the SPDF-NR and WBP-a interactions, with one neutron in the $0f_{7/2}$ and $1p_{3/2}$ levels to describe the three negative parity levels observed in the experiment.

In this work opposite-parity states are associated with configurations involving the promotion of one neutron to the full $0f_{7/2}$, $1p_{3/2}$, $0f_{5/2}$ and $1p_{1/2}$ orbitals, not only the $0f_{7/2}$ or $1p_{3/2}$ orbitals. One feature where the relevance of using the full pf orbitals for the 1p-1h excitations can be seen, is in the energy of the excited opposite-parity states. The inclusion of the full fp shell decreases the energies of the negative parity levels, moving them towards the reported experimental values.

In Table I, we show the occupancies of the ground and first excited states, and of the first excited state with opposite parity, in the ^{35,37,39,41}S isotopes, obtained with the SDPF-U interaction. The proton occupation numbers in the sd shell of the low lying states have a smooth isotopic dependence. For the ground state the occupancies of the proton $d_{5/2}$ orbital ranges between 5.34 to 5.80, while the $\pi s_{1/2}$ and $\pi d_{3/2}$ interchange their role as the number of neutron increases. They vary from 1.80 to 0.80 for the $\pi s_{1/2}$, and from 0.45 to 1.49 for the $\pi d_{3/2}$.

The increase in the number of neutrons has obviously a direct effect in the neutron occupancies. The filling of the $\nu d_{3/2}$ orbital is apparent for the three lightest isotopes, and of the $\nu f_{7/2}$ for the other three. Also the $\nu f_{7/2}$ is central in the 1p-1h excitations for the states with negative parity. While the $\nu f_{5/2}$ and $\nu p_{1/2}$ have very low occupancies, there presence plays a relevant role in obtaining the right energetics, as mentioned above.

Lets analyze the energy spectra of each particular isotope.

The calculated sequence of positive-parity levels has a clear correspondence with the experimentally observed ones in ^{35}S, Fig. 2. The energy differences of the $7/2_1^-$, $3/2_1^-$ levels from the experiment are only a few keV. The positions of the $1/2_1^-$ and $3/2_2^-$ levels are interchanged with respect to experimental ones in both calculations, and the $1/2_1^-$ level is noticeably lower than in the experiment.

Experimental data are available up to ~ 3.5 MeV for ^{37}S, shown in Fig. 3. The ground and first excited state have negative parity, and are perfectly reproduced by both calculations. The $1/2_1^-$ and $3/2_2^-$ states are interchanged with respect to the experiment, with the $3/2_2^-$ displaced to higher energies. The second excited state has positive parity, and is better adjusted by the SPDF-U interaction. Many positive parity states are predicted, but the reported $9/2_1^+$ appears too high in the calculation.



FIG. 2. Calculated and experimental level scheme of ^{35}S.



FIG. 3. Calculated and experimental level scheme of ^{37}S.



FIG. 4. Calculated and experimental level scheme of ^{39}S. The experimental data for this isotopes have been taken from LNL annual report.



FIG. 5. Calculated and experimental level scheme of ^{41}S. The experimental data for this isotopes have been taken from LNL annual report.

TABLE I. Occupancies of proton and neutron orbitals of the ground and first excited states, and of the first excited state with opposite parity, in ^{35,37,39,41}S isotopes, for SDPF-U interaction.

J^π	$\pi d_{5/2}$	$\pi s_{1/2}$	$\pi d_{3/2}$	$\nu d_{5/2}$	$\nu s_{1/2}$	$\nu d_{3/2}$	$\nu f_{7/2}$	$\nu p_{3/2}$	$\nu f_{5/2}$	$\nu p_{1/2}$
^{35}S $3/2_1^+$	5.74	1.80	0.45	5.96	1.97	3.07				
$1/2_1^+$	5.74	1.45	0.81	5.93	1.33	3.74				
$7/2_1^-$	5.43	1.34	1.22	5.71	1.61	2.67	0.93	0.06	0.01	0.0003
^{37}S $7/2_1^-$	5.80	1.68	0.52	6.00	2.00	4.00	0.95	0.04	0.01	0.000
$3/2_1^-$	5.85	1.59	0.56	6.00	2.00	4.00	0.24	0.76	0.01	0.003
$3/2_1^+$	5.60	1.32	1.08	5.94	1.93	3.13	1.59	0.28	0.09	0.04
^{39}S $7/2_1^-$	5.75	1.15	1.11	6.00	2.00	4.00	2.59	0.30	0.08	0.03
$3/2_1^-$	5.79	1.02	1.18	6.00	2.00	4.00	2.33	0.58	0.05	0.04
$3/2_1^+$	5.53	0.94	1.52	5.92	1.91	3.17	3.24	0.52	0.17	0.06
^{41}S $5/2_1^-$	5.71	0.80	1.49	6.00	2.00	4.00	4.34	0.49	0.12	0.03
$7/2_1^-$	5.69	0.84	1.46	6.00	2.00	4.00	4.38	0.43	0.15	0.04
$1/2_1^+$	5.47	0.83	1.71	5.86	1.38	3.75	5.18	0.51	0.25	0.07
$3/2_1^+$	5.42	0.85	1.72	5.89	1.86	3.24	5.00	0.64	0.27	0.08

In ^{39}S only the $7/2_1^-$, $11/2_1^-$, $15/2_1^-$ levels have been measured, and are shown in Fig. 4. The SPDF-U calculation predicts very good results for these levels. The spin and parity of the 1724 keV level has not yet been measured. It is close to the $3/2_2^-$, $1/2_1^-$ and $3/2_1^+$ levels found in the shell-model calculations. The $3/2_1^-$ level is predicted in both calculations a few keV above the $7/2^-$ the ground state.

When reaching to ^{41}S, Fig. 5, the ground state has angular momentum and parity $5/2^-$. The experimental $(7/2^-)$ level at 449 keV is predicted to be 395.

A comparison of the reduced electric $E2$ transition probabilities (in W.u.) between the experimentally available data for ^{41}S is presented in Table II. The effective charges $e^{\pi}_{\text{eff}}=1.35e$, $e^{\nu}_{\text{eff}}=0.35e$ are used for the calculation. For ^{41}S two $B(E2)$ values are reported in the experiment. The first of them is described well by the calculations. For the second both calculations predict the values 4 times larger than in the experiment.

TABLE II. Calculated $B(E2)$ values for some transitions for ^{41}S isotopes with effective charges: e^{π}_{eff}=1.35e, e^{ν}_{eff}=0.35e. The experimental γ-ray energies corresponding to these transitions are also listed.

Nucleus	$I_i^\pi \to I_f^\pi$	E_γ (keV)	Expt.	SDPF-U
^{41}S	$7/2_1^- \to 5/2_1^-$	449	14.91(6)	12.48
	$9/2_1^- \to 5/2_1^-$	904	16.57(4)	4.27

IV. CONCLUSIONS

We have interpreted the most recently available experimental data, including both positive and negative parity states, for the ^{35,37,39,41}S isotopes, performing shell model calculations with a modern interaction in the $sd-pf$ valence space. The low energy spectra and electric $E2$ transition probabilities properties were successfully reproduced, employing 0p-0h excitations for normal party states and 1p-1h for opposite parity states. The limitations of the present model were also pointed out.

ACKNOWLEDGMENTS

We would like to thank A. Poves for critical reading of the manuscript. We also thank F. Nowacki for sharing with us the SDPF-U interaction, and K. Kaneko for some useful comments. This work was supported in part by Conacyt, México, and by DGAPA, UNAM project IN103212. MJE's work is partially supported by TWAS-CONACYT grant, CONACYT project No. 155633 and by grant F2-FA-F177 of Uzbekistan Academy of Sciences. All the calculations reported in the present paper were carried out at the DGCTIC-UNAM computational facility KanBalam and at the Tochtli facility at ICN-UNAM.

---

[1] K. Wimmer, T. Kröll, R. Krücken, V. Bildstein, R. Gernhäuser, B. Bastin, N. Bree, J. Diriken, P. Van Duppen, M. Huyse, et al., Phys. Rev. Lett. **105**, 252501 (2010).

[2] T. Motobayashi, Y. Ikeda, K. Ieki, M. Inoue, N. Iwasa, T. Kikuchi, M. Kurokawa, S. Moriya, S. Ogawa, H. Murakami, et al., Phys. Lett. B **346**, 9 (1995).

[3] B. Bastin, S. Grévy, D. Sohler, O. Sorlin, Z. Dombrádi, N. L. Achouri, J. C. Angélique, F. Azaiez, D. Baiborodin, R. Borcea, et al., Phys. Rev. Lett. **99**, 022503 (2007).

[4] C. Force, S. Grévy, L. Gaudefroy, O. Sorlin, L. Cáceres, F. Rotaru, J. Mrazek, N. L. Achouri, J. C. Angélique, F. Azaiez, et al., Phys. Rev. Lett. **105**, 102501 (2010).

[5] R. Chevrier, J. M. Daugas, L. Gaudefroy, Y. Ichikawa, H. Ueno, M. Hass, H. Haas, S. Cottenier, N. Aoi, K. Asahi, et al., Phys. Rev. Lett. **108**, 162501 (2012).

[6] R. Winkler, A. Gade, T. Baugher, D. Bazin, B. A. Brown, T. Glasmacher, G. F. Grinyer, R. Meharchand, S. McDaniel, A. Ratkiewicz, and D. Weisshaar, Phys. Rev. Lett. **108**, 182501 (2012).

[7] H. T. Fortune, Phys. Rev. C **84**, 024327 (2011).

[8] G. Neyens, M. Kowalska, D. Yordanov, K. Blaum, P. Himpe, P. Lievens, S. Mallion, R. Neugart, N. Vermeulen, Y. Utsuno, et al., Phys. Rev. Lett. **94**, 022501 (2005).

[9] G. Neyens, M. Kowalska, D. Yordanov, K. Blaum, P. Himpe, P. Lievens, S. Mallion, R. Neugart, N. Vermeulen, Y. Utsuno, et al., Phys. Rev. Lett. **94**, 022501 (2005).

[10] S. Nummela, F. Nowacki, P. Baumann, E. Caurier, J. Cederkäll, S. Courtin, P. Dessagne, A. Jokinen, A. Knipper, G. Le Scornet, et al., Phys. Rev. C **64**, 054313 (2001).

[11] V. Tripathi, S. L. Tabor, P. Bender, C. R. Hoffman, S. Lee, K. Pepper, M. Perry, P. F. Mantica, J. M. Cook, J. Pereira, et al., Phys. Rev. C **77**, 034310 (2008).

[12] D. T. Yordanov, M. Kowalska, K. Blaum, M. De Rydt, K. T. Flanagan, P. Lievens, R. Neugart, G. Neyens, and H. H. Stroke, Phys. Rev. Lett. **99**, 212501 (2007).

[13] R. Kanungo, C. Nociforo, A. Prochazka, Y. Utsuno, T. Aumann, D. Boutin, D. Cortina-Gil, B. Davids, M. Diakaki, F. Farinon, et al., Phys. Lett. B **685**, 253 (2010).

[14] O. Sorlin, Z. Dombrdi, D. Sohler, F. Azaiez, J. Timr, Y. E. Penionzhkevich, F. Amorini, D. Baiborodin, A. Bauchet, F. Becker, et al., The European Physical Journal A - Hadrons and Nuclei pp. 173–178 (2004).

[15] M. De Rydt, J. M. Daugas, F. de Oliveira Santos, L. Gaudefroy, S. Grévy, D. Kameda, V. Kumar, R. Lozeva, T. J. Mertzimekis, P. Morel, et al., Phys. Rev. C **81**, 034308 (2010).

[16] R. Broda, J. Wrzesiński, A. Gadea, N. Mărginean, B. Fornal, L. Corradi, A. M. Stefanini, W. Królas, T. Pawłat, B. Szpak, et al., Phys. Rev. C **82**, 034319 (2010).

[17] S. Bhattacharyya, M. Rejmund, A. Navin, E. Caurier, F. Nowacki, A. Poves, R. Chapman, D. O'Donnell, M. Gelin, A. Hodsdon, et al., Phys. Rev. Lett. **101**, 032501 (2008).

[18] J. Ljungvall, A. Görgen, A. Obertelli, W. Korten, E. Clément, G. de France, A. Bürger, J.-P. Delaroche, A. Dewald, A. Gadea, et al., Phys. Rev. C **81**, 061301 (2010).

[19] Z. M. Wang, R. Chapman, X. Liang, F. Haas, F. Azaiez, B. R. Behera, M. Burns, E. Caurier, L. Corradi, D. Curien, et al., Phys. Rev. C **81**, 054305 (2010).

[20] Z. M. Wang, R. Chapman, F. Haas, X. Liang, F. Azaiez, B. R. Behera, M. Burns, L. Corradi, D. Curien, A. N. Deacon, et al., Phys. Rev. C **83**, 061304 (2011).

[21] L. Gaudefroy, J. M. Daugas, M. Hass, S. Grévy, C. Stodel, J. C. Thomas, L. Perrot, M. Girod, B. Rossé, J. C. Angélique, et al., Phys. Rev. Lett. **102**, 092501 (2009).

[22] L. Cáceres, D. Sohler, S. Grévy, O. Sorlin, Z. Dombrádi, B. Bastin, N. L. Achouri, J. C. Angélique, F. Azaiez, D. Baiborodin, et al., Phys. Rev. C **85**, 024311 (2012).

[23] T. R. Rodríguez and J. L. Egido, Phys. Rev. C **84**, 051307 (2011).

[24] D. Santiago-Gonzalez, I. Wiedenhöver, V. Abramkina, M. L. Avila, T. Baugher, D. Bazin, B. A. Brown, P. D. Cottle, A. Gade, T. Glasmacher, et al., Phys. Rev. C **83**, 061305 (2011).

[25] D. Sohler, Z. Dombrádi, J. Timár, O. Sorlin, F. Azaiez, F. Amorini, M. Belleguic, C. Bourgeois, C. Donzaud, J. Duprat, et al., Phys. Rev. C **66**, 054302 (2002).

[26] A. D. Davies, Ph.D. thesis, Michigan State University (2006).

[27] K. Kaneko, Y. Sun, T. Mizusaki, and M. Hasegawa, Phys. Rev. C **83**, 014320 (2011).

[28] Y. Utsuno, T. Otsuka, T. Mizusaki, and M. Honma, Phys. Rev. C **60**, 054315 (1999),

[29] P. C. Srivastava and I. Mehrotra, Journal of Physics G: Nuclear and Particle Physics **36**, 105106 (2009).

[30] P. C. Srivastava and I. Mehrotra, Physics of Atomic Nuclei **73**, 1656 (2010).

[31] P. C. Srivastava and I. Mehrotra, The European Physical Journal A - Hadrons and Nuclei **45**, 185 (2010).

[32] P. C. Srivastava and V. Kota, Physics of Atomic Nuclei **74**, 971 (2011).

[33] P. C. Srivastava, Journal of Physics G: Nuclear and Particle Physics **39**, 015102 (2012).

[34] P. C. Srivastava and I. Mehrotra, Int. J. Mod. Phys. E **21**, 1250007 (2012).

[35] P. C. Srivastava, Mod. Phys. Lett. A **277** 1250061(2012).

[36] M. Bouhelal, F. Hass, E. Caurier, F. Nowacki and A. Bouldjedri, Nucl. Phys. A **864**, 113 (2011).

[37] F. Nowacki and A. Poves, Phys. Rev. C **79**, 014310 (2009).

[38] J. Retamosa, E. Caurier, F. Nowacki, and A. Poves, Phys. Rev. C **55**, 1266 (1997).

[39] http://www.nndc.bnl.gov/nndc/ensdf.

[40] Z. M. Wang, X. Liang, R. Chapman, F. Azaiez, J. Ollier, M. J. Burns, S. Beghini, B. R. Behera, L. Corradi, D. Curien, et al., Tech. Rep., Laboratori Nazionali di Legnaro (LNL) (2007).

[41] R. Chakrabarti, S. Mukhopadhyay, Krishichayan, A. Chakraborty, A. Ghosh, S. Ray, S. S. Ghugre, A. K. Sinha, L. Chaturvedi, A. Y. Deo, et al., Phys. Rev. C **80**, 034326 (2009).

[42] P. C. Bender, S. L. Tabor, V. Tripathi, C. R. Hoffman, L. Hamilton, A. Volya, R. M. Clark, P. Fallon, A. O. Macchiavelli, et al., Phys. Rev. C **85**, 044305 (2012).